



## 核酸去除剂

货号	规格
BDAA0051	500ml

**保存:** 室温保存

**Storage:** Store at room temperature

**特点:**

安全无毒, 不氧化, 不腐蚀, 可直接接触皮肤; 快速有效去除核酸污染。

**Features:**

Safe, non - toxic, non - oxidation, non - corrosion, direct contact with the skin;

Rapid and effective removal of nucleic acid contamination.

**使用方法:**

(1) 各种台面和仪器表面的清洁: 将本产品直接喷于台面和仪器表面, 用普通吸水纸擦净, 晾干。

(2) 移液枪的清洁: 根据移液枪说明书拆卸移液枪, 将易污染核酸的部分浸放在本产品中 5-10 分钟, 再用蒸馏水彻底浸泡冲洗后, 晾干, 装回移液枪。

(3) 玻璃和塑料器皿的清洁: 将器皿浸泡在本产品中, 5-10 分钟后取出, 再用蒸馏水彻底浸泡冲洗干净后, 晾干备用。

(4) 核酸气溶胶的清洁: 当实验室有轻度的核酸气溶胶污染, 可用本产品对实验室进行全面喷洒, 待 1-2 小时后, 并配合 1:50 稀释的 84 消毒液进行地面及台面的擦拭可有效清除核酸气溶胶污染; 当实验室核酸气溶胶污染严重时, 建议处理 2-3 次, 处理完成后可检测处理效果。

**Usage:**

(1) Clean all kinds of table and instrument surface: Spray the product directly on the table and instrument surface, wipe it with ordinary absorbent paper, and dry it.

(2) Cleaning of pipette: Disassemble the pipette according to the pipette instruction, soak the part of nucleic acid that is liable to contaminate in the product for 5-10 minutes, then thoroughly soak and rinse with distilled water, dry it, and put it back into the pipette.

(3) Cleaning of glass and plastic containers: Soak the containers in this product, take them out after 5-10 minutes, thoroughly soak and rinse them with distilled water, and then dry them for later use.



(4) Nucleic acid aerosol cleaning: When there is mild Nucleic acid aerosol pollution in the laboratory, this product can be used to spray the laboratory comprehensively, after 1-2 hours, and with 1:50 diluted 84 disinfectant to wipe the floor and table surface can effectively remove Nucleic acid aerosol pollution; When the laboratory Nucleic acid aerosol pollution is serious, 2-3 times of treatment is recommended, and the treatment effect can be detected after the treatment is completed.